

KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY
RESTRICTED VETERINARY MEDICINE



Methimazole Spot-On™


For the treatment of
hyperthyroidism in cats

Your veterinarian has prescribed your cat's medication as a once-a-day topical gel. With a topical gel, the medicine is absorbed through the skin and you don't have to battle trying to give your cat a tablet!

Your "Cat Pack" contains: 1 mL Transdermal applicators

DOSE APPLICATION

1. Call your cat and place your cat comfortably next to you.
2. With applicator close at hand, place a new finger cot on thumb and index finger.
3. Remove protective cap and **slowly** push out ____ mL of gel onto one of the protected fingers as prescribed by your veterinarian. Replace cap.
4. Put the syringe down and place the cat on your lap or in a comfortable position.
5. Immediately rub the gel into the inside hairless part of the cat's ear until the gel is absorbed.
6. Carefully remove finger cots to avoid skin contact with product residue and discard.
7. Wash hands thoroughly.
8. Should a thin film remain on the cat's ear after dosing, only clean off just prior to administering next dose.



STORAGE

Store below 25 °C (room temperature). Protect from light. Replace cap between applications and store in original container.

Bayer

BAYER NEW ZEALAND LIMITED
3 Argus Place, Hillcrest, Auckland, New Zealand.
Customer Information Line 0800 446 121.

Registered pursuant to the ACVM Act 1997, No. A10271
See www.foodsafety.govt.nz for registration conditions.

84334332